

Product:	Anti-17α-Nortestosterone Antibody				
Synonyms:	Not available				
Immunogen:	17α-19-Nortestosterone-BTG				
	1 mg PAS9707	Host Species:	Ovis aries (Sheep)	Isotype:	IgG
Pack sizes: Cat. No.	10 mg PAS9556	Host Breed:	Texel	Format:	Clear liquid
		Clonality:	Polyclonal	pH:	7.4
IgG Concentration: Lot dependent. Determined @ 280nm.		Buffer: 20mM Phosphate, 150mM Sodium Chloride.			
Recommended Working Concentration*: 0.25 µg/mL *The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.		Preservative: 0.09% Sodium Azide.			
Assessment Method: Competitive ELISA.		Species Reactivity: N/A			
Method of Purification: Salt fractionation.		Storage: Can be stored for up to 3 months at +2°C - +8°C. For long term storage, aliquot and store at ≤-20 °C. Avoid repeated freeze/thaw cycles. Product should be protected from light exposure.			
Recommended The antibody is	Applications: suitable for the development of in	nmunoassays or imr	nunoaffinity purification	columns.	
Sensitivity: 10 ng/mL 17alpha-19 Nortestosterone produces 93% inhibition in a competitive ELISA employing 17α-Nortestosterone polyclonal antibody.		Target Specificity*: Data not available.			
Related Produc	cts:	E ml			

Notes & Precautions:

Antibody can be affinity purified on request. This product as supplied is intended for research applications only, not for use in therapeutic or diagnostic applications without the expressed written authorization of Randox BioReagents. A safety data sheet (SDS) can be supplied upon request. Vial should be centrifuged briefly before opening to ensure all material is removed from the vial cap.

28-Apr-22 Information correct at time of going to print

